

**Table 3. Red and blue listed plant species likely but not confirmed to occur in the Fairy Creek watershed**

| Scientific name                              | Common name                | Habitat  | Range   | Subzone <sup>a</sup>             | BC list status |
|--|----------------------------|--|---|----------------------------------|----------------|
| <i>Aphyllon pinorum</i>                      | Pine broomrape             | Moist mesic lowlands   | Rare on S Vancouver Island, S to N Idaho and NW California  | CWHmm<br>CWHxm                   | red            |
| <i>Asplenium adulterinum</i>                 | Corrupt spleenwort         | Dry to mesic walls of limestone fissures, montane and subalpine    | Rare on Vancouver Island and lower Fraser Valley; not known elsewhere in North America; Europe                              | CWHvm<br>CWHwh<br>MHmm           | blue           |
| <i>Cardamine angulata</i>                    | Angled bittercress         | Moist woods and riverbanks, lowlands                               | Rare on Haida Gwaii and Vancouver Island; N to S Alaska and S to N California   | CWHvh<br>CHWwh<br>CWHdm          | blue           |
| <i>Cephalanthera austinae</i>                | Phantom orchid             | Moist to mesic forests, lowlands                                   | Rare in the lower Fraser Valley, SE Vancouver Island and Gulf Islands; S to Idaho and California                            | CWHxm<br>CWHdm                   | red            |
| <i>Claytonia washingtoniana</i>              | Washington springbeauty    | Moist to mesic most rock outcrops and forests, lowland and montane | Rare in SW BC; S to Colorado and California   | CWHxm<br>CWHdm                   | red            |
| <i>Corallorhiza maculata var. ozettensis</i> | Ozette coralroot           | Moist to dry forests, lowland, steppe and montane                  | Frequent throughout BC esp. on the coast; south of 55 deg N, E to Newfoundland; S to Mexico; several other inland US states | CWHxm                            | blue           |
| <i>Erythronium montanum</i>                  | White glacier lily         | Moist meadows and open forests, montane and subalpine              | Rare on Vancouver Island and the adjacent mainland (Mt Waddington area); S to N Oregon                                      | CWHvm<br>CWHvh<br>CWHds<br>MHmm  | blue           |
| <i>Geum schofieldii</i>                      | Haida Gwaii avens          | Wet rock crevices, lowland, subalpine                              | Rare on Haida Gwaii and NW Vancouver Island; endemic  | CWHvh<br>MH                      | blue           |
| <i>Githopsis specularioides</i>              | Common bluecup             | Moist seepage areas on rock outcrops, lowland                      | Rare on S Vancouver Island; S to California   | CWHmm<br>CWHxm<br>MHmm           | blue           |
| <i>Mimulus sdentatus</i>                     | Tooth-leaved monkey flower | Moist forests and streambanks, lowland and lower montane           | Rare on S Vancouver Island; S to N California   | CWHvm<br>CWHvh                   | blue           |
| <i>Mitella caulescens</i>                    | Leafy mitrewort            | Wet to moist meadows and forests, lowland and montane              | Rare on S Vancouver Island and lower Fraser Valley; S to Idaho, Montana and California                                      | CWHds<br>CWHms<br>CWHxm,<br>MHmm | blue           |
| <i>Montia chamissoi</i>                      | Chamisso's montis          | Bogs, marshes and streambanks, lowland and montane                 | Rare in coastal BC; N to Alaska, E to Manitoba; S to New Mexico and California  | CWHds<br>CWHxm                   | blue           |

|                                 |                       |   |   |                                 |      |
|---------------------------------|-----------------------|---|---|---------------------------------|------|
| <i>Oxalis oregano</i>           | Redwood-sorrel        | Moist forests, lowlands   | Rare on W coast Vancouver Island and Haida Gwaii; south to California                     | CWHvm<br>CWHvh                  | blue |
| <i>Platanthera ephemerantha</i> | White-lip rein orchid | Dry forests and forest margins, lowland and montane                             | Rare in coastal BC; range includes S Vancouver Island; N to Alaska and S to California    | CWHvh                           | blue |
| <i>Polystichum setigerum</i>    | Alaska hollyfern      | Moist to mesic shady forests, rock outcrops and lava flows, lowland and montane | Rare in coastal NW BC, N to Alaska  | CWHvm<br>CWHwm<br>CWHws         | blue |
| <i>Pyrola aphylla</i>           | Leafless wintergreen  | Forests, open woods   | Mostly on mid and E side of Vancouver Island, W to W of Port Alberni and S of Thetis Lake | CWHvm<br>CWHmm<br>CWHxm         | blue |
| <i>Rubus lasiococcus</i>        | Dwarf bramble         | Mesic to moist thickets and open forests, montane and lower subalpine           | Rare in SW BC; S to N California  | CWHds<br>CWHmm<br>CWHxm<br>MHmm | blue |
| <i>Trillium hibbersonii</i>     | Hibberson's trillium  | Moist cliffs and rocks  | W coast Vancouver Island  | CWHvh                           | blue |
| <i>Viola howellii</i>           | Howell's violet       | Mesic to moist forests, lowland   | Rare on S Vancouver Island and the lower Fraser Valley, S to Oregon                       | CWHmm<br>CWHxm<br>MHmm          | red  |

a Listed subzones are those that are the same or similar to Fairy Creek; the species may occur in other subzones as well.